



□□

□□□□□4

□□□□□

2 mins to read





□□□□□□9□9□□□ □□□.....

□□
□□□
□□□□□□□□

□□□□

□□□




□□□□□□□- □□□□**good fit**□



□□□□□□□- □□□□**good fit**□

□□□



00000010000000000000000000000000000000000000

00**3-5**0



00**3-5**0

1-4000

00**1-2**0



00**1-2**0

500000000000000000000000000000000001-200

00000



00000

00



0000

- 1. Bekkali, N., et al. (2009, Apr). Infant stool form scale: development and results. J Pediatr, 154(4), 521-526.
- 2. den Hertog, J., et al. (2012, Nov). The defecation pattern of healthy term infants up to the age of 3 months. Arch Dis Child Fetal

Neonatal Ed, 97(6), F465-470.

3. Malacaman, E.E., et al. (1985, Oct). Effect of protein source and iron content of infant formula on stool characteristics. *J Pediatr Gastroenterol Nutr*, 4(5), 771-773.

4. Koopman, R., et al. (2009, Jul). Ingestion of a protein hydrolysate is accompanied by an accelerated in vivo digestion and absorption rate when compared with its intact protein. *Am J Clin Nutr*, 90(1), 106-115.

5. Vandenplas, Y. (2017, Jul). Prevention and management of cow's milk allergy in non- exclusively breastfed infants. *Nutrients*, 9(7), E731.

6. Beckingham, I.J., & Ryder, S.D. (2001, Jan). ABC of diseases of liver, pancreas, and biliary system. Investigation of liver and biliary disease. *BMJ*, 322(7277), 33-36.

7. Dallas, D.C., et al. (2012, Apr). Digestion of Protein in Premature and Term Infants. *J Nutr Disord Ther*, 2(3), 112.