□□

□□□□□4

□□□□□

2 mins to read

□□□□□

□□

□□□□□□□□9□9□□□ □□□……

□□□□□□□□

□□□□□□□□□□□□fit□

□□ - □□□□□□□□□

□□ - □□□□□□□□□

□□□□□□□□□□□□

□□□□□□□□□□

□□□□□□□ - □□□□□□□□□□□□□□□□

□□□□□□□ - □□□□□□□□□□□□□□□□

□□□□□□□□□□□□□□□

□□□□

□□□□□□□□□□□□..... □□□□□□□□□□□□□□□□□□□□

□□□

□□□

□□□□□□□□□□□□□□□□□□□

□□□

□□

□□

□□□

□□□□□□□10□□□□□□□

Neonatal Ed, 97(6), F465-470.
3. Malacaman, E.E., et al. (1985, Oct). Effect of protein source and iron content of infant formula on stool characteristics. J Pediatr Gastroenterol Nutr, 4(5), 771-773.
4. Koopman, R., et al. (2009, Jul). Ingestion of a protein hydrolysate is accompanied by an accelerated in vivo digestion and absorption rate when compared with its intact protein. Am J Clin Nutr, 90(1), 106-115.
5. Vandenplas, Y. (2017, Jul). Prevention and management of cow’s milk allergy in non- exclusively breastfed infants. Nutrients, 9(7), E731.
6. Beckingham, I.J., & Ryder, S.D. (2001, Jan). ABC of diseases of liver, pancreas, and biliary system. Investigation of liver and biliary disease. BMJ, 322(7277), 33-36.
7. Dallas, D.C., et al. (2012, Apr). Digestion of Protein in Premature and Term Infants. J Nutr Disord Ther, 2(3), 112.